arXiv:math/0510510v1 [math.GR] 24 Oct 2005

ON THE CENTER OF A COXETER GROUP

TETSUYA HOSAKA

Abstract. In this paper, we show that the center of every Coxeter group is finite and isomorphic to $(\mathbb{Z}_2)^n$ for some $n \geq 0$. Moreover, for a Coxeter system (W,S), we prove that $Z(W) = Z(W_{S \setminus \tilde{S}})$ and $Z(W_{\tilde{S}}) = 1$, where $Z(W)$ is the center of the Coxeter group W and $\tilde{S}$ is the subset of S such that the parabolic subgroup $W_{\tilde{S}}$ is the *essential parabolic subgroup* of (W,S) (i.e. $W_{\tilde{S}}$ is the minimum parabolic subgroup of finite index in (W,S)). The finiteness of the center of a Coxeter group implies that a splitting theorem holds for Coxeter groups.

1. Introduction and preliminaries

In this paper, we investigate the center of a Coxeter group. A *Coxeter group* is a group W having a presentation

$$\langle\, S \mid (st)^{m(s,t)} = 1 \text{ for } s,t \in S \,\rangle,$$

where S is a finite set and $m : S \times S \to \mathbb{N} \cup \{\infty\}$ is a function satisfying the following conditions:

(1) $m(s,t) = m(t,s)$ for each $s,t \in S$,
(2) $m(s,s) = 1$ for each $s \in S$, and
(3) $m(s,t) \geq 2$ for each $s,t \in S$ such that $s \neq t$.

The pair (W,S) is called a *Coxeter system*. Let (W,S) be a Coxeter system. For a subset $T \subset S$, W_T is defined as the subgroup of W generated by T, and called a *parabolic subgroup*. It is known that (W_T, T) is also a Coxeter system (cf. [1] and [9]). A subset $T \subset S$ is called a *spherical subset* of S, if the parabolic subgroup W_T is finite.

Date: September 9, 2005.

2000 *Mathematics Subject Classification.* 20F55.

Key words and phrases. the center of a Coxeter group, a splitting theorem for a Coxeter group.

Partly supported by the Grant-in-Aid for Young Scientists (B), The Ministry of Education, Culture, Sports, Science and Technology, Japan. (No. 15740029).

The purpose of this paper is to prove the following theorems.

Theorem 1.1. *The center of every Coxeter group is finite and isomorphic to* $(\mathbb{Z}_2)^n$ *for some* $n \geq 0$.

A Coxeter system (W, S) is said to be *irreducible*, if for any nonempty and proper subset T of S, W does not decompose as the direct product of W_T and $W_{S \setminus T}$.

Let (W, S) be a Coxeter system. Then there exists a unique decomposition $\{S_1, \dots, S_r\}$ of S such that W is the direct product of the parabolic subgroups $W_{S_1}, \dots, W_{S_r}$ and each Coxeter system (W_{S_i}, S_i) is irreducible (cf. [1] and [9]). We define $\tilde{S} = \bigcup \{S_i \,|\, W_{S_i} \text{ is infinite}\}$, and the parabolic subgroup $W_{\tilde{S}}$ is called the *essential parabolic subgroup* of (W, S). We note that $W = W_{\tilde{S}} \times W_{S \setminus \tilde{S}}$ and $W_{S \setminus \tilde{S}}$ is finite. In [6], it was proved that the essential parabolic subgroup $W_{\tilde{S}}$ is the minimum parabolic subgroup of finite index in (W, S).

We denote the center of a group G by $Z(G)$.

We also prove the following theorem in Section 2.

Theorem 1.2. *For a Coxeter system* (W, S)*,* $Z(W) = Z(W_{S \setminus \tilde{S}})$ *and* $Z(W_{\tilde{S}}) = 1$.

For an irreducible Coxeter system (W, S), if W is infinite, then $W = W_{\tilde{S}}$. Hence Theorem 1.2 implies the following.

Corollary 1.3. *For an irreducible Coxeter system* (W, S)*, if the Coxeter group* W *is infinite, then the center of* W *is trivial.*

In [8], we have obtained some splitting theorems for CAT(0) groups whose centers are finite. Theorem 1.1 and [8, Theorem 2] implies the following splitting theorem for Coxeter groups.

Corollary 1.4. *Let* (W, S) *be a Coxeter system and let* $W = W_{S_1} \times W_{S_2}$*. Suppose that the Coxeter group* W *acts geometrically on a CAT(0) space* X*. Then there exists a closed, convex,* W*-invariant, quasi-dense subspace* $X' \subset X$ *such that* X' *splits as a product* $X_1 \times X_2$ *and the action of* $W = W_{S_1} \times W_{S_2}$ *on* $X' = X_1 \times X_2$ *is the product action.*

By [2, Lemma II.6.24], we also can obtain the following corollary.

Corollary 1.5. *Suppose that a Coxeter group* $W = W_{S_1} \times W_{S_2}$ *acts geometrically on a CAT(0) space* X*. Then* W_{S_1} *and* W_{S_2} *are convex-cocompact.*

Here the definition and some properties of *convex-cocompactness* is found in [5] and [7]. We note that "geometrically finiteness" in [5] and "convex-cocompactness" in [7] coincide.

2. Proof of the main theorems

Let (W, S) be a Coxeter system and $w \in W$. A representation $w = s_1 \cdots s_l$ $(s_i \in S)$ is said to be *reduced*, if $\ell(w) = l$, where $\ell(w)$ is the minimum length of word in S which represents w.

The following lemmas are known.

Lemma 2.1 ([1])**.** *Let* (W, S) *be a Coxeter system. Suppose that* W *is finite. Then there exists a unique element* $w_0 \in W$ *of longest length, and for each* $w \in W$, $\ell(w_0 w) = \ell(w_0) - \ell(w)$. *In particular,* $w_0^2 = 1$.

Lemma 2.2 ([1], [4, Lemma 7.11])**.** *Let* (W, S) *be a Coxeter system, let* $T \subset S$ *and let* $w \in W_T$. *Then the following statements are equivalent.*

(1) W_T *is finite and* w *is the element of longest length in* W_T.
(2) $\ell(wt) < \ell(w)$ *for each* $t \in T$.

Using lemmas above, we prove the following main theorem.

Theorem 2.3. *Let* (W, S) *be a Coxeter system and let* $Z(W)$ *be the center of* W.

(1) *For each* $w \in Z(W)$, *there exists a spherical subset* T *of* S *such that* w *is the element of longest length in* W_T.
(2) $w^2 = 1$ *for any* $w \in Z(W)$.
(3) $Z(W)$ *is finite.*
(4) $Z(W)$ *is isomorphic to* $(\mathbb{Z}_2)^n$ *for some* $n \geq 0$.
(5) $Z(W) = Z(W_{S \setminus \tilde{S}})$.
(6) $Z(W_{\tilde{S}})$ *is trivial.*

Proof. Let (W, S) be a Coxeter system and let $Z(W)$ be the center of W.

(1) Let $w \in Z(W)$ and let $w = s_1 \cdots s_l$ be a reduced representation. Then $ws_1 = s_1 w$, since $w \in Z(W)$. Hence

$$ws_1 = s_1 w = s_1(s_1 s_2 \cdots s_l) = s_2 \cdots s_l.$$

Thus $\ell(ws_1) < \ell(w)$ and $w = (s_2 \cdots s_l)s_1$ is reduced. Since $w \in Z(W)$, $ws_2 = s_2 w$. Hence

$$ws_2 = s_2 w = s_2((s_2 s_3 \cdots s_l)s_1) = (s_3 \cdots s_l)s_1.$$

Thus $\ell(ws_2) < \ell(w)$ and $w = (s_3 \cdots s_l)s_1 s_2$ is reduced. By iterating the above argument, we obtain that $\ell(ws_i) < \ell(w)$ for each $i = 1, \ldots, l$. Let $T = \{s_1, \ldots, s_l\}$. By Lemma 2.2, W_T is finite and w is the element of longest length in W_T.

(2) By (1) and Lemma 2.1, we have that $w^2 = 1$ for any $w \in Z(W)$.

(3) For a spherical subset T, let w_T be the element of longest length in W_T. By (1),

$$Z(W) \subset \{w_T \,|\, T \text{ is a spherical subset of } S\},$$

which is finite, since S is finite. Hence the center $Z(W)$ is finite.

(4) We note that $w^2 = 1$ for any $w \in Z(W)$ by (2) and $vw = wv$ for any $v, w \in Z(W)$ because $Z(W)$ is the center. Thus $Z(W)$ is isomorphic to $(\mathbb{Z}_2)^n$ for some $n \geq 0$.

(5) Let $w \in Z(W)$, let $w = s_1 \cdots s_l$ be a reduced representation and let $T = \{s_1, \ldots, s_l\}$. Then w is the element of longest length in W_T by (1). Let $s \in S \setminus T$. Then $sw = ws$, since $w \in Z(W)$. Hence $sws = w$ and $s(s_1 \cdots s_l)s = s_1 \cdots s_l$. Here we note that $s \notin T = \{s_1, \cdots, s_l\}$. By Tits's theorem in [10] and [3, p.50], $ss_i = s_i s$ for each $i = 1, \ldots, l$. This means that $st = ts$ for any $t \in T$ and $s \in S \setminus T$. Hence W splits as the product $W = W_T \times W_{S \setminus T}$. Since W_T is finite, $T \subset S \setminus \tilde{S}$ by the definition of $\tilde{S}$. Hence $w \in W_T \subset W_{S \setminus \tilde{S}}$ for each $w \in Z(W)$. Thus $Z(W) \subset W_{S \setminus \tilde{S}}$. We note that $W = W_{\tilde{S}} \times W_{S \setminus \tilde{S}}$ and $Z(W) = Z(W_{\tilde{S}}) \times Z(W_{S \setminus \tilde{S}})$. Therefore $Z(W) = Z(W_{S \setminus \tilde{S}})$.

(6) We can obtain that $Z(W_{\tilde{S}})$ is trivial from (5). $\square$

References

[1] N. Bourbaki, *Groupes et Algebrès de Lie*, Chapters IV-VI, Masson, Paris, 1981.

[2] M.R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Springer-Verlag, Berlin, 1999.

[3] K.S. Brown, *Buildings*, Springer-Verlag, 1980.

[4] M.W. Davis, *Groups generated by reflections and aspherical manifolds not covered by Euclidean space*, Ann. of Math. **117** (1983), 293–324.

[5] T. Hosaka, *Limit sets of geometrically finite groups acting on Busemann spaces*, Topology Appl. **122** (2002), 565–580.

[6] T. Hosaka, *Parabolic subgroups of finite index in Coxeter groups*, J. Pure Appl. Algebra **169** (2002), 215–227.

[7] T. Hosaka, *The interior of the limit set of groups*, Houston J. Math. **30** (2004), 705–721.

[8] T. Hosaka, *On splitting theorems for CAT(0) spaces and compact geodesic spaces of non-positive curvature*, preprint.
[9] J.E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge University Press, 1990.
[10] J. Tits, *Le problème des mots dans les groupes de Coxeter*, Symposia Mathematica, vol. 1, pp. 175–185, Academic Press, London, 1969.

DEPARTMENT OF MATHEMATICS, UTSUNOMIYA UNIVERSITY, UTSUNOMIYA, 321-8505, JAPAN
E-mail address: `hosaka@cc.utsunomiya-u.ac.jp`